



Infusion Center

KENNEBUNK | ROCK ROW | TOPSHAM | PORTSMOUTH
PHONE: (207) 303-3225 | FAX: (207) 692-2473

Gemcitabine Bladder Instillation—Order Form

Patient Name	Middle	Last	DOB
--------------	--------	------	-----

Diagnosis	Diagnosis Code
-----------	----------------

Height (cm)	Actual Weight (kg)	Allergies
-------------	--------------------	-----------

***Patient will be required to have a signed consent from Urology Office prior to administration of intravesicular gemcitabine.

GEMCITABINE DOSE TO BE ADMINISTERED:

- 1000mg diluted in 50mL Normal Saline – to be prepared in a 60mL syringe

DOSING PROTOCOL – PLEASE CHECK THE APPROPRIATE GEMCITABINE DOSING OPTION BELOW:

- ☐ Induction: 1000mg/50mL Normal Saline every 7 days for 6 weeks
- ☐ Maintenance: 1000mg/50mL Normal Saline every 7 days for 3 weeks

INSTRUCTIONS FOR ADMINISTRATION:

- Instill dose (1000mg/50mL NS) into bladder
- Clamp foley catheter for 60 minutes
- After 60 minutes, drain bladder and remove foley catheter

In the event of a hypersensitivity reaction, patients will be treated according to NECS protocols for the management of infusion-room drug reactions.

Provider Name	Provider Signature
---------------	--------------------

Date	Phone	Fax	Pager
------	-------	-----	-------